

Maxpar Anti-Human TIGIT (MBSA43)-159Tb

Catalog Number, Package Size: 3159038B, 100 tests
3159038C, 25 tests

Clone: MBSA43

Other Names: VSTM3, WUCAM

Isotype: Mouse IgG1, kappa

Reactivity: Human

Tag: 159Tb

Formulation: Antibody stabilizer with 0.05% sodium azide

Storage: Store at 2–8 °C. Do not freeze.

Application: Suspension mass cytometry

Technical Information

Description: T cell Ig and ITIM domain (TIGIT) is a transmembrane glycoprotein belonging to a poliovirus receptor family of type I proteins that binds to its ligands CD155 and CD112. TIGIT is expressed on peripheral memory and regulatory (Treg) CD4+ T cells as well as NK cells and is up-regulated following activation on naive CD4+ T cells. CD155 is a high-affinity receptor for TIGIT expressed on monocytes and CD11c+ human dendritic cells (DCs). TIGIT contains an Ig-like V-type domain and an ITIM in its cytoplasmic domain, suggesting that receptor occupancy may trigger a negative signaling event. Engagement of TIGIT by CD155 on human DCs enhance the production of IL-10 while diminishing the production of IL-12p40, and a T cell-intrinsic inhibitory effect of TIGIT-dependent T cell signaling pathway was demonstrated in murine models of experimental autoimmune encephalomyelitis (EAE).

Application: The metal-tagged antibody is designed and formulated for the application of suspension mass cytometry using the Fluidigm CyTOF® suspension systems on healthy human PBMC.

Validation: Each lot of Maxpar® antibody is quality control-tested by suspension mass cytometry analysis of stained cells using appropriate positive and negative cell staining and/or activation controls.

Recommended use: Use 1 µL for up to 3×10^6 live cells in 100 µL staining volume. We recommend titrating the antibody for optimal performance for each of the desired applications. Centrifuge the stock antibody at $12,000 \times g$ for 5 min to sediment antibody aggregates.

Fixation is typically used in intracellular staining protocols or in barcoding with the Cell-ID™ 20-Plex Pd Barcoding Kit. However, fixing before antibody staining can affect epitope structure and antibody binding, with the impact varying on the type and concentration of fixative and the protocol used. It is therefore important to perform a small, preliminary antibody staining experiment, with and without fixation, using non-critical samples.

Applicable Protocols


Before using this product, refer to the instructions in the Maxpar Cell Surface Staining with Fresh Fix Protocol (400276).

References

- Bandura, D.R. et al. "Mass cytometry: technique for real time single cell multitarget immunoassay based on inductively coupled plasma time-of-flight mass spectrometry." *Analytical Chemistry* 81 (2009): 6,813–22.
- Ornatsky, O.I. et al. "Highly multiparametric analysis by mass cytometry." *Journal of Immunological Methods* 361 (2010): 1–20.
- Bengsch, B. et al. "Epigenomic-guided mass cytometry profiling reveals disease-specific features of exhausted CD8 T cells." *Immunity* 48 (2018): 1,029–45.
- Gadalla, R. et al. "Validation of CyTOF against flow cytometry for immunological studies and monitoring of human cancer clinical trials." *Frontiers in Oncology* 9 (2019): 415.

Safety

Use standard laboratory safety protocols. Read and understand the safety data sheets (SDSs) before handling chemicals. To obtain SDSs, go to fluidigm.com and search for **3000000X**.



Human PBMC stained with anti-CD8 (RPA-T8)-146Nd and anti-TIGIT (MBSA43)-159Tb. Cells shown are gated on total T cells (CD45+CD20-CD14-CD3+).

For technical support visit techsupport.fluidigm.com. | For general support visit fluidigm.com/support.

For Research Use Only. Not for use in diagnostic procedures.

Information in this publication is subject to change without notice. **Limited Use Label License:** The purchase of this Fluidigm Instrument and/or Consumable product conveys to the purchaser the limited, non-transferable right to use with only Fluidigm Consumables and/or Instruments respectively except as approved in writing by Fluidigm. **Patent and license information:** fluidigm.com/legal/notices. **Trademarks:** Fluidigm, the Fluidigm logo, Cell-ID, CyTOF, and Maxpar are trademarks and/or registered trademarks of Fluidigm Corporation in the United States and/or other countries. All other trademarks are the sole property of their respective owners. © 2020 Fluidigm Corporation. All rights reserved. 03/31/2020