

Maxpar Anti-Human CD274/PD-L1 (MIH1)-209Bi

Catalog Number, Package Size: 3209014B, 100 tests
 3209014C, 25 tests

Clone: MIH1

Other Names: B7-H1, programmed death ligand 1

Isotype: Mouse IgG1, kappa

Reactivity: Human, Dog, Mouse

Tag: 209Bi

Formulation: Antibody stabilizer with 0.05% sodium azide

Storage: Store at 2–8 °C. Do not freeze.

Application: Suspension mass cytometry

Technical Information

Description: PD-L1 (also known as CD274, B7-H1), one of the ligands for programmed death 1 (PD-1), is an immune-inhibitory receptor belonging to the CD28/cytotoxic T lymphocyte antigen 4 (CTLA-4) family. It can deliver an inhibitory signal to PD-1/B7-1-expressing T cells, resulting in immune-suppressive effects. PD-L1 is expressed on activated T cells, B cells, NK cells, DCs, macrophages, and bone marrow-derived mast cells. PD-L1 expression is also found on a wide range of human tumors such as kidney, ovarian, bladder, breast, liver, gastric, pancreatic and non-small cell lung cancer (NSCLC). Most importantly, these studies reveal that higher expression of PD-L1 may facilitate advancement of tumor stage and increase the invasion potential. PD-L1 expression can be induced by many inflammatory mediators and cytokines, of which interferon- γ (IFN- γ) is the most potent. Blockade of the PD-1/PD-L1 interaction has been shown to reverse immune exhaustion and is a critical target of current immunotherapeutics.

Application: The metal-tagged antibody is designed and formulated for the application of suspension mass cytometry using the Fluidigm CyTOF® suspension systems on healthy human PBMC.

Validation: Each lot of Maxpar® antibody is quality control-tested by suspension mass cytometry analysis of stained cells using appropriate positive and negative cell staining and/or activation controls.

Recommended use: Use 1 μ L for up to 3×10^6 live cells in 100 μ L staining volume. We recommend titrating the antibody for optimal performance for each of the desired applications. Centrifuge the stock antibody at $12,000 \times g$ for 5 min to sediment antibody aggregates.

Fixation is typically used in intracellular staining protocols or in barcoding with the Cell-ID™ 20-Plex Pd Barcoding Kit. However, fixing before antibody staining can affect epitope structure and antibody binding, with the impact varying on the type and concentration of fixative and the protocol used. It is therefore important to perform a small, preliminary antibody staining experiment, with and without fixation, using non-critical samples.

Applicable Protocols

Before using this product, refer to the instructions in the Maxpar Cell Surface Staining with Fresh Fix Protocol (400276).

References


Bandura, D.R. et al. "Mass cytometry: technique for real time single cell multitarget immunoassay based on inductively coupled plasma time-of-flight mass spectrometry." *Analytical Chemistry* 81 (2009): 6,813–22.

Ornatsky, O.I. et al. "Highly multiparametric analysis by mass cytometry." *Journal of Immunological Methods* 361 (2010): 1–20.

Boddupalli, C.S. et al. "Interlesional diversity of T cell receptors in melanoma with immune checkpoints enriched in tissue-resident memory T cells." *JCI Insight* 1 (2016): e88955.

Safety

Use standard laboratory safety protocols. Read and understand the safety data sheets (SDSs) before handling chemicals. To obtain SDSs, go to fluidigm.com and search for **300000X**.



Three-day rested (left) or PHA-stimulated human PBMC were stained with anti-CD3 (UCHT1)-170Er and anti-PD-L1 (MIH1)-209Bi antibodies. Cells shown are gated on total live CD45+ cells.

For technical support visit techsupport.fluidigm.com. | For general support visit fluidigm.com/support.

For Research Use Only. Not for use in diagnostic procedures.

Information in this publication is subject to change without notice. **Limited Use Label License:** The purchase of this Fluidigm Instrument and/or Consumable product conveys to the purchaser the limited, non-transferable right to use with only Fluidigm Consumables and/or Instruments respectively except as approved in writing by Fluidigm. **Patent and license information:** fluidigm.com/legal/notices. **Trademarks:** Fluidigm, the Fluidigm logo, Cell-ID, CyTOF, and Maxpar are trademarks and/or registered trademarks of Fluidigm Corporation in the United States and/or other countries. All other trademarks are the sole property of their respective owners. © 2020 Fluidigm Corporation. All rights reserved. 03/31/2020