

Anti-Human/Mouse CD324/E-Cadherin-158Gd

Catalog: 3158018B Clone: DECMA-1
Package size: 100 tests Isotype: Rat IgG1


Storage: Store product at 4 °C. Do not freeze. Formulation: Antibody stabilizer with 0.05% sodium azide

Cross-reactivity: Canine, Porcine

Technical Information

Validation: Each lot of conjugated antibody is quality control-tested by $\mathsf{CyTOF}^{\textcircled{R}}$ analysis of stained cells using the appropriate positive and negative cell staining and/or activation controls.

Recommended usage: The suggested use is 1 μ L for up to 3 x 10⁶ live cells in 100 μ L. It is recommended that the antibody be titrated for optimal performance for each of the desired applications.



Human MCF-7 (top) and human Jurkat cells (bottom) stained with 158Gd-anti-E-Cadherin (DECMA-1).

Description

CD324, also known as E-cadherin (Epithelial cadherin) or uvomorulin, is a member of the cadherin superfamily. E-cadherin is a calcium-dependent, transmembrane cell-cell adhesion glycoprotein composed of 4 extracellular cadherin repeats and a highly conserved cytoplasmic tail region. It functions as a cell adhesion molecule involved in development, bacterial pathogenesis and tumor invasion. E-cadherin is widely expressed in epithelial cells in the colon, uterus, liver, keratinocytes, brain, heart, muscle, kidney and pancreas as well as erythroid cells.

References

Bandura, D. R., et al. Mass Cytometry: Technique for Real Time Single Cell Multitarget Immunoassay Based on Inductively Coupled Plasma Time-of-Flight Mass Spectrometry. *Analytical Chemistry* 81 (2009): 6,813–22.

Ornatsky, O. I., et al. Highly Multiparametric Analysis by Mass Cytometry. Journal of Immunological Methods 361 (2010): 1–20.

For technical support visit fluidigm.com/support.

For Research Use Only. Not for use in diagnostic procedures.