

Anti-Vimentin-143Nd

Pathologist-Verified Clone for Imaging Mass Cytometry™

Catalog: 3143027D

Package size and concentration: 25 µg, 0.5 mg/mL

Storage: Store at 4 °C. Do not freeze.

Reactivity: Rat, Mouse, Human, Monkey

Clone: D21H3

Isotype: Rabbit IgG

Formulation: Antibody stabilizer with 0.05% sodium azide

Application: IMC-Paraffin

Technical Information


Application: The metal-tagged antibody is designed and formulated for the application of Imaging Mass Cytometry (IMC™) using the Fluidigm Hyperion™ Imaging System on formalin-fixed, paraffin-embedded (FFPE) tissue sections.

Quality control: Each lot of conjugated antibody is quality control-tested by Imaging Mass Cytometry on tissue sections.

Recommended concentration: For optimal performance it is recommended that the antibody be titrated for the desired application. Suggested initial dilution range:
 IMC-Paraffin: 1:50 to 1:200

Description

Vimentins are type III intermediate filaments found in various non-epithelial cells, especially mesenchymal cells. They are highly expressed in fibroblasts, with low expression in T and B lymphocytes and little or no expression in Burkitt's lymphoma cell lines. They are also expressed in many hormone-independent mammary carcinoma cell lines. Dynamic structural changes and spatial reorganization of vimentin in response to extracellular stimuli help to coordinate various signaling pathways. Remodeling of vimentin and other intermediate filaments is important during lymphocyte adhesion and migration through the endothelium.



Human lung (FFPE) stained with 143Nd-anti-vimentin (D21H3) at a dilution of 1:100 (red pseudocolor), 169Tm-anti-collagen I (poly) (green pseudocolor), and iridium DNA intercalator (blue pseudocolor). Heat-mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. Scale bar size = 100 µm.

References

Chang, Q. et al. "Staining of frozen and formalin-fixed, paraffin-embedded tissues with metal-labeled antibodies for imaging mass cytometry analysis." *Current Protocols in Cytometry* 82 (2017): 12.47.1–12.47.8.

Giesen, C. et al. "Highly multiplexed imaging of tumor tissues with subcellular resolution by mass cytometry." *Nature Methods* 11 (2014): 417–22.

For technical support visit <http://techsupport.fluidigm.com>. | For general support visit www.fluidigm.com/support.

For Research Use Only. Not for use in diagnostic procedures.

This product contains antibodies manufactured by and sold under license from CST™ and licensees thereof.

Information in this publication is subject to change without notice. **Safety data sheet information:** www.fluidigm.com/sds. **Patent and license information:** www.fluidigm.com/legalnotices.

Limited Use Label License: The purchase of this product conveys to the purchaser the limited, non-transferable right to use the purchased consumable or reagent only with Fluidigm Instruments and Systems. **EU's WEEE directive information:** www.fluidigm.com/compliance. Fluidigm, the Fluidigm logo, Hyperion, Imaging Mass Cytometry, and IMC are trademarks or registered trademarks of Fluidigm Corporation in the United States and/or other countries. All other trademarks are the sole property of their respective owners. © 2017 Fluidigm Corporation. All rights reserved. 10/2017